



## First record of cuckoo wasp *Trichrysis imperiosa* (Smith) (Hymenoptera, Chrysididae) from the nest of *Sceliphron coromandelicum* (Lepeletier) (Hymenoptera, Sphecidae) in India

J. Abitha<sup>1</sup>, K. Rajmohana<sup>1\*</sup>, C. Bijoy<sup>2</sup>, P. G. Aswathi<sup>2</sup> and P. Girish Kumar<sup>3</sup>

<sup>1</sup>Zoological Survey of India, M Block, New Alipore, Kolkata 700053, West Bengal, India.

<sup>2</sup>Shadpada Entomology Research Lab (SERL), Christ College (Autonomous, Affiliated to University of Calicut) Irinjalakuda 680121, Kerala, India.

<sup>3</sup>Western Ghats Regional Centre, Zoological Survey of India, Eranhipalam 673006, Kerala, India.

Email: mohana.skumar@gmail.com

**ABSTRACT:** The present study could document, *Sceliphron coromandelicum* (Lepeletier) (Hymenoptera: Sphecidae) as the host of the cuckoo wasp, *Trichrysis imperiosa* (Smith) (Hymenoptera: Chrysididae) from Kerala, India. This is the first host record of *T. imperiosa*. Interesting observations and notes on their natural history are also reported. © 2022 Association for Advancement of Entomology

**KEYWORDS:** Host association, notes, natural history, Kerala, kleptoparasite

The Chrysididae popularly called “gold wasps” or “jewel wasps” are brightly coloured and shiny Hymenoptera, mostly brilliant metallic green, violet, gold and/or red (Rosa *et al.*, 2021a). They are also termed cuckoo wasps, since they use the nest of another species for laying eggs and rearing their own young. Evolutionarily they are specialized to defend themselves during oviposition; they curl into a defensive ball through conglobulation. Their strongly chitinized and sculptured body serve to defend the attack of their hosts (Houston, 2011). The natural history of chrysidid wasps remains poorly known, though they are widespread and are important natural enemies of several groups of Hymenoptera like Sphecidae, Eumeninae and Pompilidae (Kimsey and Bohart, 1991; Bank *et al.*, 2017; Sann *et al.*, 2018). Genus *Trichrysis* Lichtenstein, 1876 of subfamily Chrysidinae are parasitoids of sphecid or crabronid wasps (Rosa *et*

*al.*, 2016) and also pompilids (Pärn *et al.*, 2015). The genus is distributed in Palaearctic, Afrotropical, Oriental and Australian Regions (Rosa *et al.*, 2016). A total of 121 species of Chrysididae under 20 genera and four subfamilies are known from India (Rosa *et al.*, 2021a; Rosa *et al.*, 2021b; Rosa and Halada, 2021; Aswathi and Bijoy, 2021).

The mud dauber wasp nest was collected from Pilassery, Kozhikode, Kerala (11.324°N; 75.9076°E) on 24-06-2021. The nest was initially kept for emergence; however, it was opened later for further studies. The fully developed individuals of both the sphecids and chrysidids were pinned and the rest were preserved in alcohol (70%). The mounted specimens were studied and photographed using Leica DFC 500 digital camera attached to Leica M205 A stereomicroscope (1X objective), and processed with LAS version 3.6, extended focus

\* Author for correspondence

software. The voucher specimens are deposited in the National Zoological Collections of Zoological Survey of India, Kolkata.

The cuckoo wasp was identified as *T. imperiosa* (Fig. 1.a), with the help of taxonomic keys (Rosa *et al.*, 2021a). With five teeth on the apex of the 3<sup>rd</sup> metasomal tergite, it belongs to *T. lusca* species group. *T. imperiosa* is similar to *T. lusca* (Fabricius, 1804), however, differs in the colour of dorsal mesosoma, in the nature of frontal carina, and also the sculpture of second and third metasomal tergites. The body is metallic greenish-blue to blue and has golden reflections on face. The species is widely distributed in India and are documented from the states of Assam, Karnataka, Kerala, Maharashtra, Meghalaya, Sikkim, West Bengal and Arunachal Pradesh (Rosa *et al.*, 2021a). Elsewhere the species is known from China (Tsuneki, 1970); Australia, Myanmar, Sri Lanka (Bingham, 1903); Vietnam (Kimsey and Bohart, 1991); Indonesia, Nepal, Papua New Guinea and Thailand (Rosa *et al.*, 2016).

Morphological identification of *S. coromandelicum* (Fig. 1. b), was made using the key to species (Anagha *et al.*, 2021). It can be easily recognized by the pronotal collar with yellowish-brown band, black metasoma with fine setae and the yellow or yellowish-brown petiole. The species is distributed in Bangladesh, Cambodia, India, Laos, Malaysia, Myanmar, Sri Lanka, Thailand, Ukraine (Pulawski, 2021; Anagha *et al.*, 2021). In India, the species is documented from Andaman Islands, Assam, Bihar, Delhi, Goa, Himachal Pradesh, Karnataka, Kerala, Maharashtra, Meghalaya, Odisha, Pondicherry, Punjab, Sikkim, Tamil Nadu, Uttarakhand, Uttar Pradesh and West Bengal (Anagha *et al.*, 2021).

**Natural history of *T. imperiosa* and *S. coromandelicum*:** The female sphecids build their mud nests in a variety of sheltered and dry sites. They are also common in human habitations (Camillo, 2002). In the present study, the mud nests of *S. coromandelicum* were found on an unplastered, illuminated wall of a building, well protected from rain and sunlight, at a height of about 1.5 m from ground level. The nest had 10 subcylindrical cells, arranged in tiers (Fig. 1c),

including an unopened cell. On breaking the single unopened cell, it was seen that the cells were provisioned with a host larva and 15 spiders, belonging to the families Clubionidae and Salticidae. Both juveniles and subadults (Fig. 1 d) could be seen. The host association of *T. imperiosa* could be ascertained since all the cells of the nest were not parasitized and three host sphecids too emerged from the same nest. In total, three *T. imperiosa* and two *S. coromandelicum* were found inside the sphecid nest (Table 1). It could be confirmed that similar to the host sphecid, only a single individual of chrysidid wasp developed from each cell. *T. imperiosa* individuals had constructed a mirror-like diaphragm across the center of the host cocoon which separated it from the host remains (Fig. 1. c).

Table 1. Details of the nest contents in each cell

No.	Cell content	Remarks
1	Vacant cell	Opened
2	Vacant cell	Opened
3	Vacant cell	Opened
4	<i>T. imperiosa</i>	Unopened
5	<i>T. imperiosa</i>	Unopened
6	<i>T. imperiosa</i>	Unopened
7	<i>T. imperiosa</i>	Unopened
8	<i>S. coromandelicum</i>	Unopened
9	<i>S. coromandelicum</i>	Unopened
10	15 spiders, <i>S. coromandelicum</i> larva	Unopened

All cuckoo wasps are parasitoids or kleptoparasites of Hymenoptera. They lay their eggs inside the host nest. In some species the hatched larva of the cuckoo wasp will consume the host larva when it is fully developed or in others it will start feeding the host egg or larvae as well as the provisioned food immediately after hatching (Szczepko *et al.*, 2003).




Fig. 1. a) *Trichrysis imperiosa*, b) *Sceliphron coromandelicum*, c) Cuckoo wasp inside *Sceliphron* nest with mirror like diaphragm, d) Spider prey and larvae

Host-parasite associations of *Trichrysis* wasps have been documented by several authors (Dufour and Perris, 1840; García Mercet, 1911; Alfken, 1915; Enslin, 1921; Trautmann, 1927; Grandi, 1931, 1936; Danks, 1971; Groot, 1971; Lomholdt, 1975; Morgan, 1984; Kimsey and Bohart, 1991; Asís *et al.*, 1994; Kunz, 1994; Strumia, 1997; Rosa, 2006). Recently Pärn *et al.*, (2015) included some Pompilidae species as potential hosts for *Trichrysis*.

*T. lusca* is reported as a parasitoid of two species of *Sceliphron* - *S. fabricator* Smith (Mocsáry, 1889, 1912; Linsenmaier, 1959) and *S. inflexus* Sickmann (Tsuneki, 1955). A few unidentified species of Eumenidae were also documented as hosts of this species (Kimsey and Bohart, 1991). The cuckoo wasps with pollen-collecting species as hosts as in the case of bees may act as parasitoids rather than kleptoparasites (Pauli *et al.*, 2019). Accordingly, in the present study since the host species *S. coromandelicum* is not a pollen collector, it can be assumed that *T. imperiosa* is in the role of a kleptoparasite rather than a parasitoid. This is the first ever host record of *T. imperiosa* globally.

### ACKNOWLEDGEMENTS

The authors are grateful to the Director, Zoological Survey of India, Kolkata for providing facilities for accomplishing this work. The authors also thank Mr. Akhilesh Mangalassery, for his wholehearted support in the field collection. First and fourth authors are grateful to the Council of Scientific and Industrial Research for the Junior Research Fellowship.

### REFERENCES

- Alfken J.D. (1915) Verzeichnis der Goldwespen (Chrysiden) Nordwestdeutschlands. *Abhandlungen des Naturwissenschaftlichen Vereins zu Bremen* 23: 291–295.
- Anagha S., Kumar P.G., Binoy C., Mazumdar P.C. and Sureshan P.M. (2021) A review of the mud-dauber wasps of genus *Sceliphron* Klug (Hymenoptera: Sphecidae) from India. *Zootaxa* 4969(1): 61–85.
- Asís J.D., Tormos J., Gayubo S.F. (1994) Biological observations on *Trypoxylon attenuatum* and description of its mature larva and its natural enemy *Trichrysis cyanea* (Hymenoptera: Sphecidae: Chrysididae). *Journal of the Kansas Entomological Society* 67: 199–200.
- Aswathi P.G. and Bijoy C. (2021) First report of the cuckoo wasp *Chrysidea falsa* (Hymenoptera) from India. *Taprobanica* 10 (2): 124–125.
- Bank S., Sann M., Mayer C., Meusemann K., Donath A., Podsiadlowski L., Kozlov A., Petersen M., Krogmann L., Meier R., Rosa P., Schmitt T., Wurdack M., Liu S., Zhou X., Misof B., Peters R.S. and Niehuis O. (2017) Transcriptome and target DNA enrichment sequence data provide new insights into the phylogeny of vespidae wasps (Hymenoptera: Aculeata: Vespidae). *Molecular Phylogenetics and Evolution* 116: 213–226.
- Bingham C.T. (1903) *The Fauna of British India, Including Ceylon and Burma. Hymenoptera, Vol. II. Ants and Cuckoo-wasps.* Taylor and Francis, London. 528 pp.
- Camillo E. (2002) The natural history of the mud dauber wasp *Sceliphron fistularium* (Hymenoptera: Sphecidae) in southeastern Brazil. *Revista de Biología Tropical* 50 (1): 127–134.
- Danks H.V. (1971) Biology of some stem-nesting aculeate Hymenoptera. *Transactions of the Royal Entomological Society* 122(11): 323–399.
- Dufour L. and Perris E. (1840) Sur les Insectes Hymenopteres qui nichent dans l'intérieur des tiges seches de la Ronce. *Annales de la Société entomologique de France* 9: 1–53.
- Enslin E. (1921) Zur Biologie des *Solenius rubicola* Duf. et Perr. (larvatus Wesm.) und seiner Parasiten. *Konowia* 1(1–2): 1–15.
- Fabricius J.C. (1804) *Systema Piezatorum secundum ordines, genera, species, adjectis synonymis, locis, observationibus, descriptionibus.* Brunsvigae. 439 pp.
- García Mercet R. (1911) Sobre la nidificación, la biología y los parásitos de algunos Esfégidos. I Congrès International d'Entomologie, Bruxelles 1: 457–464.
- Grandi G. (1931) Contributi alla conoscenza biologica e morfologica degli Imenotteri melliferi e predatori. XII. *Bollettino del Laboratorio di Entomologia di Bologna* 4: 19–71.
- Grandi G. (1936) Contributi alla conoscenza degli Imenotteri Aculeati. XVI. *Bollettino del Laboratorio di Entomologia di Bologna* 9: 253–346.

- Groot W. (1971) Waarnemingen aan Hymenoptera-nesten. *Entomologische Berichten* 31: 168–175.
- Houston T. (2011) Cuckoo Wasps (family Chrysididae). Western Australian Museum. <http://museum.wa.gov.au/research/collections/terrestrial-zoology/entomology-insectcollection/entomology-factsheets/cuckoo-wasps> (Accessed on 25 January, 2022).
- Kimsey L.S. and Bohart R.M. (1991 [1990]) *The Chrysidid Wasps of the World*. Oxford University Press, New York. 652 pp.
- Kunz P.X. (1994) *Die Goldwespen (Chrysididae) Baden-Württembergs. Taxonomie, Bestimmung, Verbreitung, Kartierung und Ökologie. Veröffentlichungen für Naturschutz und Landschaftspflege in Baden-Württemberg* 77: 188 pp.
- Lepelletier De Saint Fargeau A.L.M. (1845) *Histoire naturelle des Insectes. Hyménoptères. Vol. 3*. Librairie Encyclopédique de Roret, Paris. 646 + 4 pp.
- Lichtenstein J. (1876) Note sur le genre *Chrysis*. *Petites Nouvelles Entomologiques* 145: 27.
- Linsenmaier W. (1959) Revision der Familie Chrysididae (Hymenoptera) mit besonderer Berücksichtigung der europäischen Spezies. *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 32 (1): 1–232.
- Lomholdt O. (1975) The Sphecidae (Hymenoptera) of Fennoscandia and Denmark. *Fauna Entomologica Scandinavica* 4: 1–452.
- Mocsáry A. (1889) *Monographia Chrysididarum Orbis Terrarum Universi*. Hungarian Academy of Science, Budapest.
- Mocsáry A. (1912) *Species Chrysididarum novae. III. Annales Historico-Naturales Musei Nationalis Hungarici* 10(1): 549–592.
- Morgan D. (1984) Cuckoo-Wasps Hymenoptera, Chrysididae. *Handbooks for the Identification of British insects*. Royal Entomological Society of London 6(5): 1–37.
- Pärn M., Soon V., Vallisoo T., Hovi K., Luig J. (2015) Host specificity of the tribe Chrysidini (Hymenoptera, Chrysididae) in Estonia ascertained with trap-nesting. *European Journal of Entomology* 112(1): 91–99.
- Pauli T., Castillo Cajas, R.F., Rosa P., Kukowka S., Berg A., van den Bergh E. and Niehuis O. (2019) Phylogenetic analysis of cuckoo wasps (Hymenoptera: Chrysididae) reveals a partially artificial classification at the genus level and a species rich clade of bee parasitoids. *Systematic Entomology* 44(2): 322–335.
- Pulawski W.J. (2021) *Catalog of Sphecidae*. (Accessed on 30 January, 2022). <https://www.calacademy.org/scientists/projects/catalog-of-sphecidae>.
- Rosa P. (2006) *I Crisidi della Valle d'Aosta. Monografie del Museo regionale di Scienze naturali*, 6, St.-Pierre, Aosta. 368 pp.
- Rosa P. and Halada M. (2021) New species and new records of cuckoo wasps (Hymenoptera: Chrysididae) from India and Sri Lanka. *Zoosystematica Rossica* 30 (2): 190–212.
- Rosa P., Aswathi P.G. and Bijoy C. (2021a) An annotated and illustrated checklist of the Indian cuckoo wasps (Hymenoptera: Chrysididae). *Zootaxa* 4929(1): 1–100.
- Rosa P., Baiocchi D., Halada M. and Proshchalykin M.Y. (2021b) A new species and new records of cuckoo wasps from Pakistan and India (Hymenoptera, Chrysididae). *Journal of Hymenoptera Research* 84: 283–294.
- Rosa P., Wei N.-S., Feng J. and Xu Z.-F. (2016) Revision of the genus *Trichrysis* Lichtenstein, 1876 from China, with description of three new species (Hymenoptera, Chrysididae). *Deutsche Entomologische Zeitschrift* 63 (1): 109–136.
- Sann M., Niehuis O., Peters R.S., Christoph M., Kozlov A., Podsiadlowski L., Bank S., Meusemann K., Misof B., Bleidorn C. and Ohl M. (2018) Phylogenomic analysis of Apoidea sheds new light on the sister group of bees. *BMC Evolutionary Biology* 18 (1): 71.
- Smith F. (1874) A revision of the Hymenopterous Genera *Cleptes*, *Parnopes*, *Pyria* and *Stilbum*, with descriptions of new species of those genera, and also of new species of the Genus *Chrysis* from North China and Australia. *Transactions of the Entomological Society of London* 7: 451–471.
- Strumia F. (1996) *Praestochrysis* from India and South-East Asia (Hymenoptera Chrysididae). *Bollettino della Società entomologica italiana* 128(1): 57–64.
- Szczepko K., Kruk A., Bartos M. and Wiśniowski B. (2013) Factors influencing the diversity of cuckoo wasps (Hymenoptera: Chrysididae) in the post

- agriculture area of the Kampinos National Park, Poland. *Insect Conservation and Diversity* 6(3): 339-353.
- Tsuneki K. (1955) *Chrysis (Pentachrysis)* of North-Eastern Asia (Hymenoptera, Chrysididae). *Memoirs of the Faculty of Liberal Arts, Fukui University, Series II, Natural Science* 4(5): 35–46.
- Tsuneki K. (1970) Ein beitrag zur goldwespen-fauna Formosas. *Etizenia* 49: 1–21.
- Trautmann W. (1927) *Die Goldwespen Europas*. Uschman, Weimar. 194 pp.

*(Received July 04, 2022; revised ms accepted October 16, 2022; published December 31, 2022)*